

**Q: -How does load-following benefit native fish?
-How is it low-risk to disadvantage non-natives and not affect native fish?**

A: GCMRC believes the benefits to native fish will accrue indirectly through a reduction in predation/competition by non-native fish, primarily salmonids in the LCR reach. The model developed by Dave Speas and Carl Walters provides support for load-following to negatively impact RBT by interfering with and disrupting spawning activity as well as reducing the recruitment of young fish. The working assumption is that non-native fish overwhelm any benefits to native fish from management flows that improve habitat conditions.

Load-following destabilizes or reduces the permanence of near shoreline habitat. Conceptually, YOY for most species of fish are found along the lentic edge of the river, occupying habitat that maximizes feeding opportunities while decreasing risk from predation. When there is a change in stage fish must move laterally while following the vertical change in water edge. Repeated movement by fish and increased distance traveled, results in lateral and longitudinal displacement that increases the likelihood for predation. The greater the magnitude in change and rate the greater the effect.

Load-following can equally affect both non-native and natives fishes. There are thought to be few HBC in the mainstem during the time of the year that load-following would occur and therefore, HBC will not be negatively affected. However, the hypothesis being tested is that the current abundance of nonnative fish has created a predator/competitor load that is too excessive. It is the abundance of predators that has overwhelmed survivorship of YOY along shoreline. The intent behind this flow prescription is to disadvantage non-natives rather than benefit native fish. This experiment would test alternate hypotheses whether or not YOY survivorship is habitat limited or is predator limited.

Q: If you won't benefit native fish in the fall, why do you go to steady flows?

A: There are a number of hypotheses to test:

Ho: Turbidity reducing predation

- This hypothesis, that turbidity reduces predation by visual sight feeders is considered hierarchically second to the reduction in predator loads.

Ho: Potential benefit to native fish from more stable near shore habitat.

- GCMRC has stated that the greatest number of YOY HBC enter the mainstem during fall monsoon events, therefore there is also the potential for stable flows following these events to provide warmer and more suitable habitat conditions for these fish during the fall. GCMRC should evaluate the downstream warming potential and the increase in suitable habitat of these mainstem conditions during the Sept – Nov time period.
- Expect this potential benefit may be overcome by non-native interactions.

Ho: Lower flows result in the retention of a more natural sediment mix and higher mainstem sediment concentrations for bar building in January. The more natural mix of sediment (fines through sand) will result in better bar stability and also affect nutrient cycling

Q: How can you conduct low steady flows or low load-following flows? Don't they violate the AOP?

A: The concerns expressed about monthly volumes and the AOP process are important ones. Annual release volumes are defined by statute and resulting operating criteria. Monthly release volumes are determined to meet annual requirements, support firm power generation commitments, and address the combined risks of powerplant bypasses, and over- and under-release of annual volumes. The GCPA did not alter the water development philosophy of the previous 1956 and 1968 Acts; thus, the financial integrity of the Basin Fund is of great concern to the States. As the AOP is developed each year, consideration is given to all the factors listed in the 1970 Operating Criteria, which include environmental concerns. The issue raised on the Feb. 8 conference call was not so much that the monthly volumes cannot be adjusted, but rather that the process of developing those monthly volumes must remain consistent with the water development and financial goals of previous Acts. If monthly volumes need to be adjusted from the typical decision making process in order to conduct an experiment, this issue would need to be addressed during the preparation of the 2003 AOP.

Q: If this is an 8.23 MAF year, July and August would be 800,000 Acre Feet and September, October, and November would be low volume months. How can you conduct the flows you describe, given these potential volumes?

A: The proposed hydrograph shows both the end of the 2002 water year (expected to be 8.23 maf) and the entire 2003 water year (expected to be about 10 maf). No adjustment to the 2002 water year hydrograph is proposed, with the possible exception of reacting to fall tributary inputs. Thus, July and August of the 2002 water year are expected to be about 800,000 af months, and GCMRC will monitor the effects of ROD releases during this year as part of a "baseline" data collection to determine the effects of ROD operations.

The reaction to fall tributary inputs is to recommend either a reduction to low steady releases (below 10,000 cfs) to conserve sediment inputs in the main channel or low fluctuating flows (5,000 – 10,000 cfs) to conserve sediment inputs in the eddies, or lastly the release of a HMF.

Q: Could you use a HMF to conserve sediment instead of low steady flows or low load-following flows?

A: You might potentially store some sand and possibly some limited amount of finer sediment within eddies, but the total sand conservation achieved would be far less than if the entire channel bed was available for storage of fines under the 10,000 cfs or less flow alternative. A comparison of results from the 1996 BHBF, and the 1997, 31,500 cfs spike flow provide some valuable insights on this matter. One important concept to keep in mind is that the total potential storage for sand within eddies is only a small fraction of what can be stored within the main channel, under flow conditions that limit downstream transport.

Eddy-sand bars studied following the November 1997, spike flow (under relatively sediment-supply enriched conditions), showed a much finer grain-size distribution than bars created by the 1996 BHBF (under relatively greater sediment-depleted conditions). The 1997 eddy deposits were relatively cohesive compared with the 1996 deposits, and were much darker in color,

suggesting some higher content of organics. Unfortunately, the average thickness of most of the 1997 deposits was on the order of 10-20 cm, or relatively thin compared with bar thickness measured after the 1996 BHBF. One preliminary conclusion derived from the 1997, sediment experiment was that the 31,500 cfs spike flow produced a “stage-limited” response, and that thicker and larger bars might have been deposited in November 1997, had the stage been increased to above peak power plant levels. Flows preceding the November 1997 test were not as low as those being proposed for future experiments, yet there was still evidence to suggest that at least some portion of the summer sediment inputs remained in the channel by the time the high flow occurred.

Q: Could UC and LC talk about a way to work this out if deliveries are lower and water volumes need to be shifted for this experiment, especially since releases from Mead wouldn't be affected?

A: This can be done as long as the Compact and the specific annual release requirements contained in the 1970 Operating Criteria can't be violated.

Q: Are the purposes responsive to the motion? Materials describe two purposes: (1) sediment, and (2) benefits native fish. I thought the motion was primarily about sediment.

A: The point of the motion was primarily to test the sediment Ad hoc group's second recommendation, but GCMRC interpreted the motion as a sediment conservation experiment within the framework of benefiting native fish.

Q: Is the GCMRC draft proposal in concert with the RPA?

A: Yes. As per the FWS RPA, the draft proposal identifies the need for the development of experimental flows to test, or the implementation of alternate flows as specified in the Biological Opinion of GCD. One such flow was the 2000-LSSF experiment (SWCA 2000) that used a holistic hydrograph that was intended to produce habitat improvements that would benefit HBC, rather than a mechanistic approach specifically applied to a life history component. The approach here is more discrete and testable than the former flow scenario.

Q: Would load-following confound measuring impacts of Brown Trout removal in Bright Angel?

A: Load-following is designed to be a more systematic treatment. One year of load following vs. one-year of Brown Trout evaluations are very different. Having both would enhance the potential effort.

The hypothesis that is being tested is whether or not predation is limiting YOY HBC recruitment. Whether we have multiple treatments that are each reducing predator loads does not affect the overall nature of the test. For the test, it is not what reduces trout more effectively (flows, tributary treatments, mechanical removal), but whether the reduction of trout abundance by some

means has an effect on YOY HBC recruitment? Depending on the ultimate test response (which remains unknown), a titration approach could be implemented that would allow for a measure of efficiency of one method over the use of an alternate method.

Q: Are the wheels in motion to do Brown Trout removal this Fall?

A: Grand Canyon National Park is supportive but the resources (staff and funding) are not there to do it this Fall.

Q: If FY 2002 is an 8.23 MAF year and one needs to take information from Summer of 2000 and compare it to ROD flows in 2001 and 2002, do you have the baseline data collection for that comparison?

A: A similar level of effort for monitoring downstream native and non-native fish has continued since the LSSF flows of 2000. Seining at a lower level of effort is continuing. The integrated sediment data is continuing to be collected. Reduced efforts on downstream temperature is ongoing. The level of resolution and many of the specific studies intended to answer specific effects of the LSSF treatment have not been continued. Once a set of experimental flows is established and the hypotheses to be tested are articulated, GCMRC will need to review the power of the existing monitoring activities.

Q: Is there any response of HBC to ROD flow changes? Do we know if there are changes in the LCR that may be the cause and that will be ignored by this effort?

A: We are beginning to look at the LCR hydrology record. One would expect to see changes in LCR hydrology due to development in the watershed resulting in groundwater depletion, surface water retention, and changes runoff patterns in response to paved areas, among others. It is not clear if we will be able to correlate these with changes in HBC. The working hypothesis is that predation and competition from non-natives will overwhelm habitat effects. This hypothesis also assumes that some portion of the LCR population recruited from fish which spent part of their early life history in the mainstem and that this portion of the recruitment is being negatively affected by non-native fishes.

There are other hypotheses that might explain the decline in abundance of HBC. Although there is considerable uncertainty around all of these alternate hypotheses. Not all of the hypotheses are experimentally testable to explain the causal nature of the recent decline of HBC. For this reason, we are suggesting an experiment to reduce predators in the mainstem because it is a practical and low risk treatment. However, if we were to implement such an experiment it does not imply that the current effort monitoring the species of concern would be discontinued, or that alternate mechanisms or factors that might be also contributing to the decline would be ignored, unconsidered or omitted from the present monitoring.

Q: Are there other options for sediment conservation in September-December? Options might include HMFs or steady 8,000 -10,000 cfs flows? Load-following between 5000-9000 cfs? Would the conceptual model or Wiele efforts help sort this out?

A: The conceptual model is not predictive in the sense that we can calculate the expected outcome of an experiment, but it may have utility in comparing the results of several options for portions of the proposed hydrograph.

Wiele's bar-evolution modeling simulations indicate that bars are most effectively deposited when a combination of high suspended-sediment concentrations and high-stage conditions occur simultaneously. Limiting downstream transport of newly input fine sediment until the release of a controlled flood is the most probable means of achieving both of the above conditions, if releasing a controlled flood during or immediately after tributary inputs is not a viable option. The conceptual model's sediment dynamic sub-model is based on the same basic premise: when abundant sand supply is available, then it will get mobilized by a high-flow release and the result will be an increase in sand storage at high stage elevations.

Q: What reservoir elevation do we need to be at to use the spillways?

A: By January 2003, Lake Powell storage elevation needs to be at 3657 feet to have 14,000 cfs of flow in either of the spillways. This level of discharge is needed for the spillway to function in a safe manner. For January 2003, the lake level is projected to be 3648 feet. As a result, there seems to be little hope that the BHBF proposed for January 2003 could exceed 45,000 cfs.

Q: December is a high power demand month, is it possible to do fluctuating flows up to 15,000 cfs. What would this do to sediment storage for a January BHBF?

A: Load following between 9,000 and 15,000 cfs during December 2002, is certainly one of the possibilities being evaluated following summer/fall sediment inputs. If the sediment inputs occurred in September, and the flows were released to 10,000 cfs or less through November, then there would be less downstream transport during the proposed December load following than if normal operations occurred throughout the September through December period. The impact on downstream transport of newly input sand during the December operations is most easily evaluated by reviewing figure 2, included in the Rubin et al. memorandum of August 2000.

Assuming: 1) that at least 500,000 tons of sand entered the main channel from the Paria River in September 2002, 2) that flows were immediately reduced to 10,000 cfs or less from the time of the input until December 1st, and 3) that the average flow for December 2002, was about 12,000 cfs – figure 2 suggests that about one-half of the newly input sediment introduced in September 2002, would be transported downstream in about 45 to 250 days, once December's operations started. Likewise, if we assume that the proposed BHBF occurred at the start of January 2003, then there is reason to believe with some certainty that less than one-half of the September sediment input would be exported downstream before the January BHBF was released.

Q: If sediment is stored in the eddies in an HMF, why do you go to monthly low flows?

A: The HMF approach foregoes hypotheses related to storing the fuller range of sediment components and the turbidity effects. Also, there is only a limited volume of sand that can be stored in the eddies and it is much less than the volume of fine-sediment that can be stored in the channel bed throughout the entire river channel. Results of the November 1997, sediment experiment (31,500 cfs) indicated that sand storage increased within eddies, but that this increase in storage was very limited compared to the volume of sand that might have been conserved had a higher flow been released that would have taken advantage of higher-elevation storage locations along shorelines. There is no basis at this time for concluding that the potential eddy-storage volume within the ecosystem is sufficient to sustain sand-bar maintenance long term. In light of this, the most conservative scientific experimental approach is to conduct the flow experiment that has the greatest likelihood of optimizing sediment conservation and sand-bar restoration/maintenance. Doing less optimal sediment treatments in the future may very well be justified during periods when reduced-flow operations are not possible, or on the basis of new findings that indicate that the fullest range of channel storage is not required for long-term sand bar sustainability.

Q: Why not test a BHBF in Fall?

A: This is viewed as outside the current interpretation of the Law of the River.

Q: Could you move the load-following period to February to address the possible public concern of stranding adult trout?

A: This can be considered. One is not sure if it will have the same effect of reducing non-native spawning. Using slower down ramp strategies as well could reduce stranding of adults.

Q: Concerned that load-following is outside the ROD? Do we have compliance for fluctuating flows that violate daily ranges and proposed upramp rates?

A: GCMRC believes that this would be covered as an experiment under the ROD. We do need to check into compliance.

Q: I want to ask you to look at trade-off in fine sediment losses through a two-stage approach: -HMF followed by a BHBF and the benefit of doing low load-following during winter-peak demand

A: Because we don't have a very robust numerical predictive capability, with respect to sediment dynamics in this system, the best way of quantifying these differences is by conducting the proposed experiment, and then comparing the results to those of the proposed alternatives. The next best way is by using empirically derived methods for estimating sediment transport, such as the method shown in figure 2, of the Rubin et al. (2000) memorandum.

Q: Given that the motion focuses on sediment, how does the low load-following benefit the sediment?

A: The winter load following component of the experiment is not intended to benefit the sediment resource. However, there may be some advantage derived from the winter load-following with respect to modifying the newly formed bar morphologies.

Q: Will the load-following erode the recently deposited sediment from the BHBF? What sediment-related hypotheses does the load-following test?

A: Any operation following the BHBF will result in reworking of the newly deposited sand bars. This bar reworking can be viewed as “erosion,” but there is no way to avoid it. The basic question remains as to what operation will export the least volume of sand downstream for any range of antecedent sand-storage conditions? A secondary consideration concerns how the operations that follow the BHBF will affect the morphology of the new bars with respect to ecosystem value. A few hypotheses might be considered:

Ho: Winter load-following does not result in a increased export of stored sediment.

Ho: Winter load-following does not modify new eddy bars in a manner that makes them more stable of greater resource value.

Ho: Winter load-following does not achieve eddy bar morphologies that allow for greater access to recreational users.

Q: There are non-ROD elements in Figure 1, load-following, low Fall releases, could include BHBF outside ROD period and HMF outside the ROD. Recommends two Figures: an ideal hydrograph and one that balances legal/policy trade-offs.

A: Given the provision in the ROD that calls for experiments if the resources are not responding as expected to the ROD flows, no elements of the proposed flows are thought to be outside the ROD. GCMRC and the TWG should review this more thoroughly.

Q: Why is a fall BHBF outside Law of the River?

A: The current interpretation of the Law of the River would prevent a BHBF from occurring for other than dam safety purposes. The triggering criteria developed to address that provide a window between January to July for conducting BHBFs.

Q: Why isn't this a program of flows?

A: Given that the motion asks for a proposal to be brought back to the AMWG in April for WY 2002-2003, this is intended to be an element of a program of flows. It is clear from the conference call that there are a number of alternative ideas for experimental flows. This is the flow GCMRC recommends the AMP try first, with the understanding that this single year's flow would fit into an overall program of flows.

Q: Integrated Ecosystem Experiment: I recommend that the experiment should be less ambitious and should simply test the sediment paradigm. The experiment should be within Law of the River and the ROD. We do have to get through July and August with meeting power supplies.

A: The AMP is intended to use an ecosystem-science approach in testing the effects of dam operations. It also recognizes that the Endangered Species Act mandates that the Department of the Interior take action to protect these species. At the January AMWG meeting, GCMRC presented data that indicated sediment resources and HBC are not responding to the ROD as anticipated in the EIS. It would be irresponsible for GCMRC to make a recommendation to the AMWG for an experimental flow that does not take an ecosystem science approach and that does not address the resources of concern. It is our belief that the flows proposed in the February 8 (Version 1.1) draft are within that portion of the ROD that allows for experimentation, if ROD flows are not achieving the intended benefits. GCMRC also believes that after the AOP process considers the experimental flow request and determines WY 2003 monthly volumes, the proposed flows are in compliance with the current interpretation of the Law of the River as all of the flow elements that are proposed for testing could be implemented, following appropriate compliance, within the current interpretation of the Law of the River.

Q: If you go to the low load-following proposed, can you test the up- and down-ramp rates and their effects?

A: Addressing the sediment-transport dynamic issues related to ramping rates is a difficult challenge that requires very high-resolution data sets, or a very sophisticated numerical predictive capability. GCMRC has been examining an optically-based technique (LISST) for measuring suspended sediment. Results to date look promising. We will know by summer if we can use this approach to examine changes on up- and down-ramp rates. In the event that LISST technologies prove adequate for monitoring suspended-sediment transport in the Colorado River ecosystem, then these methods will be used for evaluating issues of ramping rates and relationships to sand resources, etc. Researchers at University of Arizona have studied issues of ramping rates and their impacts on the dynamics of sand bars previously. These EIS results can be reviewed again, and additional numerical simulations can be run and evaluated, pending approval by managers and on the basis of funding availability.

Q: What's next? When will the public be informed?

A: This is the conceptual phase. The questions and concerns we heard today will be used to refine what we bring to the TWG in February. The TWG will provide an opportunity for public input. Following the TWG, a recommendation will be forwarded to the AMWG. We assume that detailed design work with scientists and detailed public impact will occur following that and final approval will be obtained in July? GCMRC and the AGFD held a preliminary meeting with the Fishing Guides at Lees Ferry on February 12 to provide the same conceptual material that was presented on February 8 and invite their input.

Q: In your next draft, please address impact this might have on the old high water zone (OHWZ) vegetation?

A: Okay.

Q: How much money do we have for this?

A: The experimental flow fund will contain about \$1 million by January 2003. Any remaining funding needs will be sought as an appropriation request or through reprioritizing AMP activities.